

# Exact Simulation of Stochastic Volatility and other Affine Jump Diffusion Processes

Mark Broadie

Columbia University, Graduate School of Business, 415 Uris Hall, 3022 Broadway, New York, NY,  
10027-6902, mnb2@columbia.edu

Özgür Kaya

Columbia University, Department of Industrial Engineering and Operations Research, 500 West 120th  
Street, New York, NY, 10027-6699, ok94@columbia.edu

First draft: July 23, 2003

This version: September 29, 2004

## Abstract

The stochastic differential equations for affine jump diffusion models do not yield exact solutions that can be directly simulated. Discretization methods can be used for simulating security prices under these models. However, discretization introduces bias into the simulation results and a large number of time steps may be needed to reduce the discretization bias to an acceptable level. This paper suggests a method for the exact simulation of the stock price and variance under Heston's stochastic volatility model and other affine jump diffusion processes. The sample stock price and variance from the exact distribution can then be used to generate an unbiased estimator of the price of a derivative security. We compare our method with the more conventional Euler discretization method and demonstrate the faster convergence rate of the error in our method. Specifically, our method achieves an  $O(s^{-\frac{1}{2}})$  convergence rate, where  $s$  is the total computational budget. The convergence rate for the Euler discretization method is  $O(s^{-\frac{1}{3}})$  or slower, depending on the model coefficients and option payoff function.

*Subject Classifications:* Simulation, efficiency: exact methods. Finance, asset pricing: computational methods.

**Acknowledgement:** This paper was presented at seminars at Columbia University, the sixth Monte Carlo and Quasi-Monte Carlo Methods in Scientific Computing conference, and INFORMS Annual Meeting in Atlanta. We thank seminar participants and an anonymous referee for helpful comments and suggestions. This work was supported in part by NSF grant DMS-0074637.

# 1 Introduction

Financial models usually specify the dynamics of the state variables, e.g., stock price, volatility and interest rate, as stochastic differential equations (SDE). If these SDEs yield closed form solutions, then Monte Carlo simulation can be used to generate an unbiased estimator of the price of a derivative security. When using Monte Carlo simulation, many sample paths of the state variables are generated and the payoff of the derivative is evaluated for each path. Discounting and averaging over all paths gives an estimator of the derivative price. The error in the Monte Carlo estimator can be calculated using the central limit theorem and converges to zero as the number of sample paths used increases. If generating each sample path requires roughly equal amount of time, then the convergence rate for such an unbiased Monte Carlo estimator is  $O(s^{-\frac{1}{2}})$  where  $s$  is the total computational budget.

If the SDEs that define the dynamics of the state variables do not yield closed form solutions, it is still possible to use Monte Carlo simulation by discretizing the time interval and simulating the state process dynamics on this discrete time grid. However, the approximation of continuous time processes by discrete time processes introduces bias into the simulation estimator. This bias causes several important problems when estimating the prices or greeks of derivative securities. First, since the magnitude of the bias is unknown, it is difficult to obtain valid confidence intervals. Second, many time steps may be necessary to reduce the bias to an acceptable level, and even more computational effort is needed to verify that the bias is small enough. Finally, both the number of time steps and number of sample paths need to be increased together in order to decrease the total error of the simulation estimator, but the optimal choice of these parameters is difficult to specify in advance. Duffie and Glynn [12] study optimal rules for allocating the total computational budget between the number of time steps and the number of simulation trials. They show that, under some regularity conditions, the error for a first order method such as Euler discretization has  $O(s^{-\frac{1}{3}})$  convergence,

In this paper, we propose a method that recovers the  $O(s^{-\frac{1}{2}})$  convergence rate of an unbiased Monte Carlo estimator for simulating derivative prices under some affine jump diffusion models. We first consider the stochastic volatility (SV) model of Heston [19], which models the variance as a square-root process that is correlated with the stock price. Scott [29] uses Fourier inversion methods to sample from the integral of a square-root process in an interest rate setting. Willard [31] observes that the stock price is lognormally distributed conditional on a path of the Brownian motion driving the variance

process. A similar ‘mixing result’ was independently discovered in Romano and Touzi [28]. We build on the ideas from these papers to generate an exact sample from the distribution of the terminal stock price. We first generate a sample from the final value of the variance. Then using Fourier inversion methods and conditioning, we get a sample from the integral of the variance. Finally, conditional on the values for the variance process, we are able to generate a sample from the stock price. We also show how to extend the method for models involving jumps in addition to the stochastic volatility. Strong empirical evidence of stochastic volatility and jumps in financial markets is documented in many recent papers, including Bates [6], Bakshi et al. [4], Chernov et al. [9], Eraker et al. [15], Pan [26] and others.

Some other authors used Monte Carlo simulation for pricing derivatives under the stochastic volatility models. When the stock price and volatility are instantaneously uncorrelated, Hull and White [20] show that the stock price has a lognormal distribution conditional on the integral of the variance process. They write the price of a call option as a Black-Scholes price given the integral of the variance. They simulate paths of the variance process and find the unconditional price of the option by taking the average of the conditional Black-Scholes prices. Willard [31] considers the case when the stock price and volatility are instantaneously correlated. He observes that the stock price is lognormally distributed conditional on the entire path of the Brownian motion driving the variance process, and the call option price can be written as a Black-Scholes price conditional on this path. He simulates the variance process on a discrete time grid and uses conditional Monte Carlo to get the option price. His method still suffers from discretization bias because of the discretization of the variance process. Heath and Platen [18] propose a variance reduction technique based on Itô calculus and show how their method can be applied for simulation under Heston’s stochastic volatility model. Their method uses discretization methods to simulate the state processes, and therefore does not eliminate the discretization bias.

The rest of the paper is organized as follows: Section 2 introduces the SV model dynamics and Euler discretization method. Section 3 is a step by step introduction to our method for exact simulation. In Section 4, we give some numerical results and compare our method to Euler discretization. In Section 5, we show that the efficiency of our simulation method can be further improved using conditional Monte Carlo techniques. Section 6 shows how to extend the method to models that involve jumps. In Section 7, we give an application of the exact simulation algorithm to the pricing of forward start options. The derivation of the characteristic function of the integral of the variance is given in the Appendix.

## 2 SV Model and Euler Discretization

### 2.1 Specification of the SV Model

Heston's model [19] is based on the following equations, which represent the dynamics of the stock price and the variance processes under the risk-neutral measure:

$$dS_t = rS_t dt + \sqrt{V_t} S_t \left[ \rho dW_t^{(1)} + \sqrt{1 - \rho^2} dW_t^{(2)} \right] \quad (1)$$

$$dV_t = \kappa(\theta - V_t) dt + \sigma_v \sqrt{V_t} dW_t^{(1)}. \quad (2)$$

The first equation gives the dynamics of the stock price:  $S_t$  denotes the stock price at time  $t$ ,  $r$  is the risk neutral drift,  $\sqrt{V_t}$  is the volatility. The second equation gives the evolution of the variance which follows the square-root process:  $\theta$  is the long-run mean variance,  $\kappa$  represents the speed of mean reversion, and  $\sigma_v$  is a parameter which determines the volatility of the variance process.  $W_t^{(1)}$  and  $W_t^{(2)}$  are two independent Brownian motion processes, and  $\rho$  represents the instantaneous correlation between the return process and the volatility process.

### 2.2 Euler Discretization

Euler discretization can be used to approximate the paths of the stock price and variance processes on a discrete time grid. Let  $[0 = t_0 < t_1 < \dots < t_M = T]$  be a partition of time interval into  $M$  equal segments of length  $\Delta t$ , i.e,  $t_i = iT/M$  for each  $i = 0, 1, \dots, M$ . The discretization for the stock price process is:

$$S_{t_i} = S_{t_{i-1}} + rS_{t_{i-1}}\Delta t + \sqrt{V_{t_{i-1}}} S_{t_{i-1}} \left[ \rho \Delta W_{t_i}^{(1)} + \sqrt{1 - \rho^2} \Delta W_{t_i}^{(2)} \right] \quad (3)$$

where  $\Delta W_{t_i}^{(j)} = W_{t_i}^{(j)} - W_{t_{i-1}}^{(j)}$ ,  $j = 1, 2$ . The discretization for variance process is:

$$V_{t_i} = V_{t_{i-1}} + \kappa(\theta - V_{t_{i-1}})\Delta t + \sqrt{V_{t_{i-1}}} \sigma_v \Delta W_{t_i}^{(1)}. \quad (4)$$

To simulate the Brownian increments, the fact that each increment  $W_{t_i}^{(j)} - W_{t_{i-1}}^{(j)}$  is independent of others is used. Each such increment is normally distributed with mean 0 and standard deviation  $\sqrt{\Delta t}$ . In particular, we can simulate these increments by first generating a uniform random variate

$U$  and then replacing  $\Delta W_{t_i}^{(j)}$  by  $\sqrt{\Delta t}F^{-1}(U)$ , where  $F(x)$  is the cumulative distribution function of a standard normal random variable. There are efficient algorithms for accurately approximating the inverse function of a normal distribution function. We use the algorithm by Moro [25]. For each increment, we use an independent uniform random variate, so we need a total of  $2M$  uniform random variates to simulate a path of stock price and variance processes. When using Euler discretization with a large number of simulation trials and a large number of time steps, the number of uniform random variates needed may be huge. We use the random number generator by Matsumoto and Nishimura [24], which has a very long period. To avoid the negative values for variance and stock price, we set these to zero if we encounter negative values during the simulation.

By repeating this procedure, many paths can be generated. Because no-arbitrage derivative prices are given by their expected discounted payoffs under the risk-neutral measure, the price of a derivative, written as  $E[f(S_T)]$ , can be estimated by Monte Carlo simulation using:

$$\frac{1}{N} \sum_{k=1}^N f(\hat{S}_T^M) \quad (5)$$

where  $N$  is the number of sample paths used in simulation and  $\hat{S}_T^M$  denotes the simulated value of  $S_T$  over each sample path using  $M$  time steps. This Monte Carlo estimator converges to the correct price  $E[f(S_T)]$  as the number of time steps  $M$  and the number of samples  $N$  become large.

There are two types of error associated with the Monte Carlo estimator given in (5). First there is a statistical error which decreases at the rate  $O(1/\sqrt{N})$ . This is due to the central limit theorem: the standard error of the simulation estimator is  $\sigma_f/\sqrt{N}$  where  $\sigma_f$  is the standard deviation of  $f(\hat{S}_T^M)$ . Second, there is a discretization error or discretization bias defined as:  $|E[f(S_T)] - E[f(\hat{S}_T^M)]|$ . This error is caused by simulation of continuous-time processes using discrete-time approximations. Kloeden and Platen [22] show that, under the conditions stated in their Theorem 14.5.2, this error decreases at the rate  $O(1/M)$  for Euler discretization. Talay and Tubaro [30], and Bally and Talay [5] give similar results, and show that the error can be expanded in powers of  $1/M$  under some regularity conditions on the coefficients of the SDEs and the function  $f$ . The conditions for first order convergence do not hold for the SDEs in (1) and (2), so the actual convergence may be slower in this case.

Duffie and Glynn [12] study the optimal allocation of a computational budget between the number

of time steps and number of simulation trials. Their result implies that it is optimal to choose the number of time steps proportional to the square-root of the number of simulation trials for first order methods, although the optimal constant of proportionality is left undetermined. They show that with their allocation rule, the convergence rate of the error for the Euler discretization is  $O(s^{-\frac{1}{3}})$  where  $s$  is the total computational budget. More generally, if  $p$  is the convergence order of the bias, then an optimal allocation has error convergence of  $O(s^{-\frac{p}{2p+1}})$ . However, since the conditions for first order convergence do not hold,  $O(s^{-\frac{1}{3}})$  convergence rate is not guaranteed in this case. Indeed, the numerical examples we consider in Section 4 demonstrate that Euler discretization may achieve a lower convergence rate depending on the convergence rate of the bias.

There are higher order discretization schemes that can be used instead of Euler for discretization of stock price and variance processes. For example, the Milstein scheme is a discretization scheme that achieves second order convergence for the bias under the conditions given in Kloeden and Platen [22]. In theory, such a second order method would improve the convergence of the total simulation error to  $O(s^{-\frac{2}{5}})$ . But again, since the conditions for convergence are not satisfied by the dynamics of the state variables, this method is not guaranteed to achieve a second order convergence rate. Also, the convergence is not very smooth. For an example of this for the SV model, see Glasserman [17]. Another disadvantage of the Milstein scheme is that it is harder to implement and takes more time per replication than the Euler discretization.

### 3 Exact Simulation of the SV Model

In this section we give the details of the exact simulation method. In particular, we show how to generate an exact sample from the distribution of  $S_t$  by conditioning on the values generated by the variance process.

The stock price at time  $t$ , given the values of  $S_u$  and  $V_u$  for  $u < t$ , can be written as:

$$S_t = S_u \exp \left[ r(t-u) - \frac{1}{2} \int_u^t V_s ds + \rho \int_u^t \sqrt{V_s} dW_s^{(1)} + \sqrt{1-\rho^2} \int_u^t \sqrt{V_s} dW_s^{(2)} \right] \quad (6)$$

and the variance at time  $t$  is given by the equation:

$$V_t = V_u + \kappa\theta(t - u) - \kappa \int_u^t V_s ds + \sigma_v \int_u^t \sqrt{V_s} dW_s^{(1)}. \quad (7)$$

We go through the following steps to sample from the distribution of  $(S_t, V_t)$ :

Exact Simulation Algorithm for the SV Model:

1. Generate a sample from the distribution of  $V_t$  given  $V_u$ .
2. Generate a sample from the distribution of  $\int_u^t V_s ds$  given  $V_t$  and  $V_u$ .
3. Recover  $\int_u^t \sqrt{V_s} dW_s^{(1)}$  from (7) given  $V_t$ ,  $V_u$  and  $\int_u^t V_s ds$ .
4. Generate a sample from the distribution of  $S_t$  given  $\int_u^t \sqrt{V_s} dW_s^{(1)}$  and  $\int_u^t V_s ds$ .

Next we specify the details of each of these steps.

### 3.1 Sampling from $V_t$ given $V_u$

The distribution of  $V_t$  given  $V_u$  for some  $u < t$  is up to a scale factor, a noncentral chi-squared distribution as noted by Cox et al. [11]. The transition law of  $V_t$  can be expressed as:

$$V_t = \frac{\sigma_v^2(1 - e^{-\kappa(t-u)})}{4\kappa} \chi_d'^2 \left( \frac{4\kappa e^{-\kappa(t-u)}}{\sigma_v^2(1 - e^{-\kappa(t-u)})} V_u \right), t > u \quad (8)$$

where  $\chi_d'^2(\lambda)$  denotes the noncentral chi-squared random variable with  $d$  degrees of freedom, and noncentrality parameter  $\lambda$ , and

$$d = \frac{4\theta\kappa}{\sigma_v^2}. \quad (9)$$

This says that, given  $V_u$ ,  $V_t$  is distributed as  $\sigma_v^2(1 - e^{-\kappa(t-u)})/(4\kappa)$  times a noncentral chi-squared random variable with  $d$  degrees of freedom and noncentrality parameter

$$\lambda = \frac{4\kappa e^{-\kappa(t-u)}}{\sigma_v^2(1 - e^{-\kappa(t-u)})} V_u. \quad (10)$$

Thus, we can sample from the distribution of  $V_t$  exactly, provided that we can sample from the noncentral chi-squared distribution.

Let  $\chi_d^2$  denote a chi-squared random variable with  $d$  degrees of freedom. Johnson et al. [21] show

that for  $d > 1$ , the following representation is valid:

$$\chi_d'^2(\lambda) = \chi_1'^2(\lambda) + \chi_{d-1}^2. \quad (11)$$

To generate  $\chi_d'^2(\lambda)$ ,  $d > 1$ , we can generate  $\chi_{d-1}^2$  and an independent standard normal random variable  $Z$  and set

$$\chi_d'^2(\lambda) = (Z + \sqrt{\lambda})^2 + \chi_{d-1}^2. \quad (12)$$

Thus, sampling from a noncentral chi-squared distribution is reduced to sampling from an ordinary chi-squared and an independent normal when  $d > 1$ .

For any  $d > 0$ , a noncentral chi-squared random variable can be represented as an ordinary chi-squared random variable with a random degrees of freedom parameter. In particular, if  $N$  is a Poisson random variable with mean  $\frac{1}{2}\lambda$ , then  $\chi_{d+2N}^2$  has the same distribution as  $\chi_d'^2(\lambda)$ . So, to sample from  $\chi_d'^2(\lambda)$ , we can first generate a Poisson random variable  $N$  and then, conditional on  $N$ , we can sample a chi-squared random variable with  $d + 2N$  degrees of freedom.

In the simulation of  $V_t$ , we use the representation given in (12) when  $d > 1$ , and we use the Poisson method described above otherwise. To sample from the chi-squared distribution, the methods to sample from gamma distribution can be used since chi-squared is a special case of this distribution. We use algorithms GS\* and GKM3 in Fishman [16] to sample from the gamma distribution. The first of these is used when the degrees of freedom parameter for chi-squared distribution is less than one, and the second one is used otherwise.

### 3.2 Sampling from $\int_u^t V_s ds$ given $V_t$ and $V_u$

Once we have a sample for  $V_t$ , we want to sample from the distribution of  $\int_u^t V_s ds$  given  $V_t$  and  $V_u$ . Scott [29] uses Fourier inversion techniques to invert the characteristic function of this distribution to sample from the same distribution in an interest rate setting. We follow a similar approach to generate a sample for the integral. The Laplace transform

$$E \left[ \exp \left( -a^* \int_u^t V_s ds \right) \middle| V_u, V_t \right]$$

can be derived using the results in Pitman and Yor [27]. Details of the derivation are given in the appendix. The characteristic function follows by setting  $a^* = -ia$ :

$$\begin{aligned}
\Phi(a) &= E \left[ \exp \left( ia \int_u^t V_s ds \right) \middle| V_u, V_t \right] \\
&= \frac{\gamma(a) e^{\frac{-1}{2}(\gamma(a)-\kappa)(t-u)} (1 - e^{-\kappa(t-u)})}{\kappa(1 - e^{-\gamma(a)(t-u)})} \\
&\quad \times \exp \left\{ \frac{V_u + V_t}{\sigma^2} \left[ \frac{\kappa(1 + e^{-\kappa(t-u)})}{1 - e^{-\kappa(t-u)}} - \frac{\gamma(a)(1 + e^{-\gamma(a)(t-u)})}{1 - e^{-\gamma(a)(t-u)}} \right] \right\} \\
&\quad \times \frac{I_{0.5d-1} \left[ \sqrt{V_u V_t} \frac{4\gamma(a)e^{-0.5\gamma(a)(t-u)}}{\sigma^2(1 - e^{-\gamma(a)(t-u)})} \right]}{I_{0.5d-1} \left[ \sqrt{V_u V_t} \frac{4\kappa e^{-0.5\kappa(t-u)}}{\sigma^2(1 - e^{-\kappa(t-u)})} \right]} \tag{13}
\end{aligned}$$

where  $\gamma(a) = \sqrt{\kappa^2 - 2\sigma^2 ia}$ ,  $d$  is as given in (9) and  $I_\nu(x)$  is the modified Bessel function of the first kind.

The probability function can be computed using Fourier inversion methods. If we let  $V(u, t)$  denote the random variable that has the same distribution as the integral  $\int_u^t V_s ds$  conditional on  $V_u$  and  $V_t$ , then we can write (see Feller [14]):

$$F(x) \equiv \Pr\{V(u, t) \leq x\} = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\sin ux}{u} \Phi(u) du = \frac{2}{\pi} \int_0^{\infty} \frac{\sin ux}{u} \text{Re}[\Phi(u)] du. \tag{14}$$

We use the trapezoidal rule to compute the probability distribution numerically:

$$\Pr\{V(u, t) \leq x\} = \frac{hx}{\pi} + \frac{2}{\pi} \sum_{j=1}^{\infty} \frac{\sin hjx}{j} \text{Re}[\Phi(hj)] - e_d(h) \tag{15}$$

where  $h$  is a grid size to be set to achieve any desired level of accuracy, and  $e_d(h)$  is the resulting discretization error. The trapezoidal rule works well for periodic and other oscillating integrands, because the errors tend to cancel. The discretization error  $e_d(h)$  can be bounded above and below by using Poisson summation formula as shown in Abate and Whitt [1]:

$$0 \leq e_d(h) = \sum_{k=1}^{\infty} \left[ F \left[ \frac{2k\pi}{h} + x \right] - F \left[ \frac{2k\pi}{h} - x \right] \right] \leq F^c \left[ \frac{2\pi}{h} - x \right] \tag{16}$$

where  $F^c(x) = 1 - F(x)$ .

If we want to achieve a discretization error  $\epsilon$ , then the step size should be:

$$h = \frac{2\pi}{x + u_\epsilon} \geq \frac{\pi}{u_\epsilon}, \text{ where } F^c(u_\epsilon) = \epsilon \text{ and } 0 \leq x \leq u_\epsilon. \quad (17)$$

Finding the correct  $u_\epsilon$  is not straightforward but the moments of the distribution can easily be found using the characteristic function in (13). We use the mean plus five or more standard deviations to get a value with a small tail probability.

To be able to calculate  $\Pr\{V(u, t) \leq x\}$  using (15), we need to determine a point at which the summation can be terminated. Let  $N$  represent the last term to be calculated so that the approximation becomes

$$\Pr\{V(u, t) \leq x\} = \frac{hx}{\pi} + \frac{2}{\pi} \sum_{j=1}^N \frac{\sin hjx}{j} \text{Re}[\Phi(hj)] - e_d(h) - e_T(N) \quad (18)$$

where  $e_T(N)$  denotes the truncation error caused by using  $N$  terms in the finite sum. The magnitude of the characteristic function,  $|\Phi(u)|$ , has value one at  $u = 0$  and it decreases as  $u$  increases. Since  $|\sin(ux)| \leq 1$ , The integrand in (14) is bounded by:

$$\frac{2|\text{Re}[\Phi(u)]|}{\pi u} \leq \frac{2|\Phi(u)|}{\pi u} = \beta(u). \quad (19)$$

Since the integrand is oscillating, the bound for the last term gives a good estimate for the truncation error, i.e. we set  $e_T(N) = h\beta(Nh)$ . The summation is terminated at  $j = N$  when

$$\frac{|\Phi(hN)|}{N} < \frac{\pi\epsilon}{2} \quad (20)$$

where  $\epsilon$  is the desired truncation error.

It should be noted that very often the error bounds stated above for determination of the grid size  $h$  and the truncation point  $N$  are conservative. By trial and error one can get  $h$  and  $N$  that achieves the desired accuracy with fewer number of terms.

The hardest and numerically most time consuming part of (15) is the evaluation of the characteristic function. The characteristic function given in (13) involves two modified Bessel functions of the first kind, and the one in the numerator has a complex argument. The modified bessel function of the first

kind is characterized by the following power series:

$$I_\nu(z) = \left(\frac{1}{2}z\right)^\nu \sum_{j=0}^{\infty} \frac{(\frac{1}{4}z^2)^j}{j!\Gamma(\nu+j+1)} \quad (21)$$

where  $\Gamma(x)$  is the gamma function and  $z$  is a complex number. We used a routine written by Amos [3] to calculate these functions. The routine uses the power series for small  $|z|$ , the asymptotic expansion for large  $|z|$ , the Miller Algorithm normalized by the Wronskian and a Neumann Series for intermediate magnitudes, and the uniform asymptotic expansion for large orders. The routine can also calculate a scaled version of the Bessel function,  $e^{-|Re(z)|}I_\nu(z)$  which makes it possible to compute the function value for very large orders.

The representation of  $I_\nu(z)$  given in (21) is not complete because the function  $z^\nu$ , which is a factor on the right hand side, is a multivalued function and needs precise specification. It is defined to be  $\exp(\nu \log z)$ , where the argument of  $z$  is given its principal value so that

$$-\pi < \arg(z) \leq \pi.$$

This is the definition most software packages and the programming library routines uses to calculate the function value. However, the function  $I_\nu(z)$  defined in this way is discontinuous at each point along the negative  $x$ -axis as are all functions defined on a specific branch of  $\arg(z)$ . So to get around this problem and accomplish the continuity of the characteristic function, we need to keep track of  $\arg(z)$  and change the branch when necessary. We use the following continuation formula from Abramowitz and Stegun [2] for this purpose which makes it possible to calculate the value of  $I_\nu(z)$  for  $\arg(z)$  different than its principal value:

$$I_\nu(ze^{m\pi i}) = e^{m\nu\pi i}I_\nu(z) \quad \{m \text{ integer}\}. \quad (22)$$

As this formula shows, to calculate the function on a different branch, we need to multiply the value on the principal branch by a factor  $e^{m\nu\pi i}$ .

To simulate the value of the integral, we use the inverse transform method. We generate a uniform random variable  $U$  and then find the value of  $x$  for which the  $\Pr\{V(u, t) \leq x\}$  is equal to  $U$ . We use a second order Newton's method to find the solution for  $x$ . This method is used to solve the equations of

the form  $f(x) = 0$  using the iteration:

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)} \left( 1 - \sqrt{\frac{2f(x_n)f''(x_n)}{f'(x_n)^2}} \right).$$

The conditional distribution of the integral is approximately normal and the initial guess for  $x$  is calculated using the inverse normal distribution function with the mean and standard deviation of the correct distribution. The moments of the correct distribution are found using (13). We set the starting value to a small value such as 0.01 times the mean if the normal value is negative. The first and second derivatives of the distribution function are calculated numerically using the derivatives of the approximation (15). The iterative procedure achieves five decimal place accuracy in 3 to 4 iterations. We use a bisection search method when the above mentioned method fails to converge to true value, which typically occurs when  $U$  is very close to 0 or 1.

### 3.3 Generating a Sample for $S_t$

After having generated samples from  $V_t$  and  $\int_u^t V_s ds$  we use (7) to get:

$$\int_u^t \sqrt{V_s} dW_s^{(1)} = \left( \frac{1}{\sigma_v} \right) \left( V_t - V_u - \kappa\theta(t-u) + \kappa \int_u^t V_s ds \right). \quad (23)$$

Furthermore, since the process for  $V_t$  is independent of the Brownian motion  $W_t^{(2)}$ , the distribution of  $\int_u^t \sqrt{V_s} dW_s^{(2)}$  given the path generated by  $V_t$  is normal with mean 0 and variance  $\int_u^t V_s ds$ .

Using these two results, we get that the conditional distribution of  $\log S_t$  is normal with mean

$$m(u, t) = \log S_u + \left[ r(t-u) - \frac{1}{2} \int_u^t V_s ds + \rho \int_u^t \sqrt{V_s} dW_s^{(1)} \right]$$

and variance

$$\sigma^2(u, t) = (1 - \rho^2) \int_u^t V_s ds.$$

To generate a sample from  $S_t$ , we generate a standard normal random variable  $Z$  and then set:

$$S_t = e^{m(u,t) + \sigma(u,t)Z}.$$

We can use the value of  $S_t$  generated in this way to find the price of any derivative that depends on the value of  $S_t$ . Repeating over many sample paths, taking the average and discounting gives us an unbiased estimator of the derivative price.

## 4 Numerical Results

In this section we present some numerical comparisons of the Euler and the exact simulation method described above. We use a European call option for this purpose. Heston [19] gives a closed form solution for the price of this option, so we are able to calculate and compare the two methods through their root-mean-squared (RMS) errors. If  $\hat{\alpha}$  is the simulation estimator used for the derivative price and  $\alpha$  is the true price, then the bias of the estimator is given by  $(E[\hat{\alpha}] - \alpha)$  and the variance is given by  $E[(\hat{\alpha} - E[\hat{\alpha}])^2]$ . RMS error is then defined as  $(\text{bias}^2 + \text{variance})^{1/2}$ . The bias for Euler discretization with a specific number of time steps can be estimated using a very large number of simulation trials to estimate  $E[\hat{\alpha}]$ , and then taking the difference with the true price. The variance for each simulation experiment is estimated by the sample variance of the simulation output.

The simulation experiments in this and the other sections of the paper were performed on a desktop PC with an AMD Athlon 1.66 GhZ processor and 624 MB of RAM, running Windows XP Professional. The codes were written in C programming language, and the compiler used was Microsoft Visual C++ 6.0.

Table 1 gives simulation results for a European call option. The set of parameters for the SV model are taken from Duffie et al. [13]. These were found by minimizing the mean squared errors for market option prices for S&P 500 on November 2, 1993. The bias column is estimated using 40 million simulation runs. The number of time steps for the Euler discretization is set equal to the square-root of the number of simulation trials.

The time needed for the exact method is more than the Euler discretization for small number of simulation runs. However, the exact method requires less computation time as the desired accuracy is increased.

There are some cases for which the bias of the Euler discretization is very large even if a large number of time steps is used. This is especially true when  $(2\kappa\theta/\sigma_v^2) \leq 1$  and  $\sigma_v$  is large relative to  $\theta$ . In this case the variance process for Euler discretization often hits to 0 causing a large discretization bias.

Table 1: Simulation Results under the SV Model for a European Call Option

a. Simulation with the Exact Method

| No of Simulation Trials | RMS Error | Computing Time (sec.) |
|-------------------------|-----------|-----------------------|
| 10000                   | 0.0750    | 3.8                   |
| 40000                   | 0.0373    | 15.2                  |
| 160000                  | 0.0186    | 60.0                  |
| 640000                  | 0.0093    | 239.4                 |
| 2560000                 | 0.0046    | 955.7                 |
| 10240000                | 0.0023    | 3822.6                |

b. Simulation with the Euler Discretization

| No of Simulation Trials | No of Time Steps | Bias   | Standard Error | RMS Error | Computing Time (sec.) |
|-------------------------|------------------|--------|----------------|-----------|-----------------------|
| 10000                   | 100              | 0.1543 | 0.0772         | 0.1725    | 0.2                   |
| 40000                   | 200              | 0.1003 | 0.0381         | 0.1073    | 1.9                   |
| 160000                  | 400              | 0.0662 | 0.0189         | 0.0689    | 15.2                  |
| 640000                  | 800              | 0.0395 | 0.0094         | 0.0406    | 121.3                 |
| 2560000                 | 1600             | 0.0267 | 0.0047         | 0.0272    | 970.0                 |
| 10240000                | 3200             | 0.0161 | 0.0023         | 0.0163    | 7758.6                |

Option parameters:  $S = 100$ ,  $K = 100$ ,  $V_0 = 0.010201$ ,  $\kappa = 6.21$ ,  $\theta = 0.019$ ,  $\sigma_v = 0.61$ ,  $\rho = -0.70$ ,  $r = 3.19\%$ ,  $T = 1.0$  year, true option price=6.8061.

Table 2 demonstrates this case. The bias column is estimated using 40 million simulation runs.

As seen from the simulation results in Table 2, significant bias remains in the Euler discretization even when 3200 time steps are used with 10,240,000 simulation runs. The RMS error for our exact method with the same number of trials is about 30 times smaller with much less computation time. Figures 1 and 2 show the convergence of the RMS graphically for both methods. The faster convergence rate of the exact method is demonstrated by the higher slope in the graphs.

One apparent problem when using Euler discretization is the choice of the number of time steps to be used in the simulation. Our choice of setting number of the time steps equal to the square-root of the number of simulation trials is somewhat arbitrary. Duffie and Glynn [12] show that, asymptotically, it is optimal to increase the number of time steps proportional to the square-root of the number of simulation trials for first-order methods such as Euler discretization. But the optimal constant of proportionality is not easy to determine. Ideally, for a specific number of simulation trials, one should choose the number of time steps such that the magnitude of the discretization bias and the standard error will be close. Then, as more accuracy is needed, the number of time steps should be increased proportional to the square-root of the number of simulation trials. This way, simulation estimate converges to the true




Figure 1: Convergence of the RMS Errors of the Two Methods for the Option in Table 1 (SV Model)




Figure 2: Convergence of the RMS Errors of the Two Methods for the Option in Table 2 (SV Model)

Table 2: Simulation Results under the SV Model for a European Call Option

a. Simulation with the Exact Method

| No of Simulation Trials | RMS Error | Computing Time (sec.) |
|-------------------------|-----------|-----------------------|
| 10000                   | 0.6125    | 3.8                   |
| 40000                   | 0.2904    | 15.3                  |
| 160000                  | 0.1464    | 61.3                  |
| 640000                  | 0.0726    | 244.5                 |
| 2560000                 | 0.0362    | 978.6                 |
| 10240000                | 0.0181    | 3916.5                |

b. Simulation with the Euler discretization

| No of Simulation Trials | No of Time Steps | Bias   | Standard Error | RMS Error | Computing Time (sec.) |
|-------------------------|------------------|--------|----------------|-----------|-----------------------|
| 10000                   | 100              | 2.1962 | 0.6568         | 2.2923    | 0.2                   |
| 40000                   | 200              | 1.6413 | 0.3247         | 1.6731    | 1.9                   |
| 160000                  | 400              | 1.2151 | 0.1544         | 1.2248    | 15.4                  |
| 640000                  | 800              | 0.9265 | 0.0761         | 0.9296    | 122.6                 |
| 2560000                 | 1600             | 0.7093 | 0.0379         | 0.7103    | 980.4                 |
| 10240000                | 3200             | 0.5367 | 0.0188         | 0.5370    | 7838.4                |

Option parameters:  $S = 100, K = 100, V_0 = 0.09, \kappa = 2.00, \theta = 0.09, \sigma_v = 1.00, \rho = -0.30, r = 5.00\%, T = 5.0$  years, true option price=34.9998.

value with neither of the errors dominating the other. Of course, in practice, we do not usually know the true value of the quantity we want to estimate and therefore it is not possible to know what the bias is. Even for the cases where we do know the true value, we might need to do a convergence study with many simulation trials to estimate the bias. Thus, the choice of the number of time steps is another shortcoming of the Euler discretization method.

Table 3 shows the bias values for various time steps for the two options considered before. The bias values are estimated using 40 million simulation runs. For both options, the bias dominates the standard error when number of time steps,  $M$ , is set to the square-root of the number of simulation trials,  $N$ . For the first option, the number of time steps that makes bias and standard error approximately equal is between  $\sqrt{N}$  and  $10\sqrt{N}$ , while for the second option this value is greater than  $10\sqrt{N}$ .

Careful examination of the bias values in Table 3 reveals another problem with the convergence of the Euler discretization. The bias values do not decrease to half of their values when the number of time steps are doubled. We mentioned in section 2.2 that the conditions for first order convergence do not hold for the SDEs in (1) and (2). These examples demonstrate that the convergence for discretization bias may indeed be significantly slower. As shown in Duffie and Glynn [12], if  $p$  is the convergence order

Table 3: Effect of the Number of Time Steps in Euler discretization

a. Bias Values for the Option in Table 1

| No of Simulation Trials ( $N$ ) | Standard Error | Bias Values for Various Time Steps ( $M$ ) |        |                |        |                  |        |
|---------------------------------|----------------|--|--------|----------------|--------|------------------|--------|
|                                 |                | $M = 0.1\sqrt{N}$                          | Bias   | $M = \sqrt{N}$ | Bias   | $M = 10\sqrt{N}$ | Bias   |
| 10000                           | 0.0772         | 10   | 0.5348 | 100            | 0.1543 | 1000             | 0.0364 |
| 40000                           | 0.0381         | 20   | 0.3897 | 200            | 0.1003 | 2000             | 0.0225 |
| 160000                          | 0.0189         | 40   | 0.2667 | 400            | 0.0662 | 4000             | 0.0163 |
| 640000                          | 0.0094         | 80   | 0.1745 | 800            | 0.0395 | 8000             | 0.0099 |
| 2560000                         | 0.0047         | 160  | 0.1156 | 1600           | 0.0267 | 16000            | 0.0067 |
| 10240000                        | 0.0023         | 320  | 0.0739 | 3200           | 0.0161 | 32000            | 0.0029 |

a. Bias Values for the Option in Table 2

| No of Simulation Trials ( $N$ ) | Standard Error | Bias Values for Various Time Steps ( $M$ ) |        |                |        |                  |        |
|---------------------------------|----------------|--|--------|----------------|--------|------------------|--------|
|                                 |                | $M = 0.1\sqrt{N}$                          | Bias   | $M = \sqrt{N}$ | Bias   | $M = 10\sqrt{N}$ | Bias   |
| 10000                           | 0.6568         | 10   | 6.0489 | 100            | 2.1962 | 1000             | 0.8408 |
| 40000                           | 0.3247         | 20   | 4.5423 | 200            | 1.6413 | 2000             | 0.6257 |
| 160000                          | 0.1544         | 40   | 3.2997 | 400            | 1.2151 | 4000             | 0.5142 |
| 640000                          | 0.0761         | 80   | 2.4358 | 800            | 0.9265 | 8000             | 0.3731 |
| 2560000                         | 0.0379         | 160  | 1.7914 | 1600           | 0.7093 | 16000            | 0.2935 |
| 10240000                        | 0.0188         | 320  | 1.3501 | 3200           | 0.5367 | 32000            | 0.2273 |

of the bias, then an optimal allocation has error convergence of  $O(s^{-\frac{p}{2p+1}})$ . The convergence rate of the bias for these examples is close to  $O(1/\sqrt{M})$ . Assuming this rate is the real convergence rate, the convergence rate of the total error for an optimal allocation would be  $O(s^{-\frac{1}{4}})$ . In practical cases, a non-optimal allocation may be used since the real convergence rate of the bias is unknown, leading to even slower convergence rates. These arguments provide another motivation for using an exact simulation method instead of biased methods.

## 5 Conditional Monte Carlo for the SV Model

Willard [31] proposed a conditional Monte Carlo method to improve the efficiency of the simulation estimators under stochastic volatility models. His method is applicable for path-independent derivatives that have closed form solutions under the Black-Scholes model. In this section we apply the conditional Monte Carlo method to improve the efficiency of the exact simulation method we described in Section 3. We also compare the improved results with the results of Euler discretization in the same setting.

We follow the approach of Willard and use a European call option to demonstrate the conditional Monte Carlo method. We assume the dynamics of the stock price and the variance processes are as

given in (1) and (2). Let  $BS(S_0, \sigma)$  denote the Black-Scholes formula for a European call option with maturity  $T$ , strike  $K$  and written on a stock with initial price  $S_0$  and constant volatility  $\sigma$ . Let  $\tilde{\sigma}$  be the average volatility of the stock over the time horizon:

$$\tilde{\sigma} \equiv \sqrt{\frac{1}{T} \int_0^T V_s ds}. \quad (24)$$

Willard observes that conditional on a path of the Brownian motion driving the variance process, the price of the call option can be written as

$$BS(S_0 \xi, \tilde{\sigma} \sqrt{1 - \rho^2}) \quad (25)$$

where

$$\xi = \exp \left( -\frac{\rho^2}{2} \int_0^T V_s ds + \rho \int_0^T \sqrt{V_s} dW_s^{(1)} \right). \quad (26)$$

Then, using the law of iterated expectations, the unconditional price of the option can be written as:

$$E \left[ BS \left( S_0 \xi, \tilde{\sigma} \sqrt{1 - \rho^2} \mid W_s^{(1)} \right) \right]. \quad (27)$$

The notation used emphasizes that the Black-Scholes price inside the expectation is conditional on a path of the Brownian motion  $W_s^{(1)}$ . Actually, it is not necessary to condition on the entire path of the Brownian motion. We can just write:

$$E \left[ BS \left( S_0 \xi, \tilde{\sigma} \sqrt{1 - \rho^2} \mid \int_0^T V_s ds, \int_0^T \sqrt{V_s} dW_s^{(1)} \right) \right] \quad (28)$$

since we only need  $\int_0^T V_s ds$  and  $\int_0^T \sqrt{V_s} dW_s^{(1)}$  to compute the Black-Scholes price. We can now use the above representation to estimate the option price using conditional Monte Carlo.

The above approach can also be used to generate unbiased estimators of the Greeks under the SV model. For example, to calculate the delta of a European call option, we can differentiate the expression in (28) and get:

$$E \left[ \Delta \left( S_0 \xi, \tilde{\sigma} \sqrt{1 - \rho^2} \mid \int_0^T V_s ds, \int_0^T \sqrt{V_s} dW_s^{(1)} \right) \xi \right] \quad (29)$$

where  $\Delta(S, v)$  denotes the Black-Scholes delta. See Broadie and Glasserman [7] for more details about

the pathwise method to estimate option greeks in a Monte Carlo simulation. Extensions of the exact simulation method in this paper to the development of unbiased simulation estimates of Greeks for various exotic options are given in Broadie and Kaya [8].

In Section 3, we described the steps to sample from the final variance value  $V_T$ , and the integral of the variance value  $\int_0^T V_s dW_s^{(1)}$ , and also showed how to recover  $\int_0^T \sqrt{V_s} dW_s^{(1)}$  given these two. When using the exact simulation method with conditional Monte Carlo, we can complete these steps and then use (25) to get the conditional option price. By repeating over many paths and taking the average, we can get an unbiased estimator of the option price.

When using Euler discretization with conditional Monte Carlo, only the variance process needs to be simulated using the discretization in (4). Simulation of the stock price is not needed, because after simulating a path of the variance process, one can use (25) to find the conditional option price. Using conditional Monte Carlo cuts the simulation time for Euler discretization roughly in half since only one process is simulated instead of two. However, it does not eliminate the discretization bias because the variance process is still being simulated on a discrete time grid.

Tables 4 and 5 show the simulation results for the two options considered in the previous section, this time using conditional Monte Carlo with the exact simulation and Euler discretization methods. Although the conditional Monte Carlo decreases the simulation time and the standard error for the Euler discretization, the bias is almost unchanged. Figures 3 and 4 show the convergence of the RMS errors.

Comparing the RMS error for the exact method using conditional Monte Carlo and unconditional Monte Carlo, we see that conditional Monte Carlo can lead to significant variance reduction. For the second option considered, more than 50-fold variance reduction is achieved using conditional Monte Carlo.

In addition to the conditional Monte Carlo method described above, Willard [31] uses quasi-Monte Carlo (low-discrepancy) methods, and shows that it results in significant efficiency improvements. Quasi-Monte Carlo helps in improving the convergence of the simulation error, but it does not decrease or eliminate the discretization bias. Therefore, qualitative results similar to the ones above will hold in comparing the exact method and the Euler discretization method when quasi-Monte Carlo is used for both.




Figure 3: Convergence of the RMS Errors of the Two Methods for the Option in Table 4 (SV Model)




Figure 4: Convergence of the RMS Errors of the Two Methods for the Option in Table 5 (SV Model)

Table 4: Simulation Results for a European Call Option using Conditional Monte Carlo

a. Simulation with the Exact Method

| No of Simulation Trials | RMS Error | Computing Time (sec.) |
|-------------------------|-----------|-----------------------|
| 10000                   | 0.0395    | 3.6                   |
| 40000                   | 0.0199    | 14.5                  |
| 160000                  | 0.0099    | 57.7                  |
| 640000                  | 0.0050    | 230.4                 |
| 2560000                 | 0.0025    | 921.6                 |
| 10240000                | 0.0012    | 3687.4                |

b. Simulation with the Euler discretization

| No of Simulation Trials | No of Time Steps | Bias   | Standard Error | RMS Error | Computing Time (sec.) |
|-------------------------|------------------|--------|----------------|-----------|-----------------------|
| 10000                   | 100              | 0.1574 | 0.0421         | 0.1630    | 0.1                   |
| 40000                   | 200              | 0.1000 | 0.0207         | 0.1022    | 1.0                   |
| 160000                  | 400              | 0.0649 | 0.0102         | 0.0657    | 8.1                   |
| 640000                  | 800              | 0.0419 | 0.0050         | 0.0422    | 64.1                  |
| 2560000                 | 1600             | 0.0267 | 0.0025         | 0.0268    | 511.1                 |
| 10240000                | 3200             | 0.0170 | 0.0012         | 0.0170    | 4082.8                |

Option parameters:  $S = 100$ ,  $K = 100$ ,  $V_0 = 0.010201$ ,  $\kappa = 6.21$ ,  $\theta = 0.019$ ,  $\sigma_v = 0.61$ ,  $\rho = -0.70$ ,  $r = 3.19\%$ ,  $T = 1.0$  year, true option price=6.8061.

## 6 Extensions to Other Models

### 6.1 Simulation Under the SVJ Model

The stochastic volatility with jumps (SVJ) model is an extension of SV model to include jumps in the stock price process. The risk-neutral dynamics are:

$$dS_t = (r - \lambda\bar{\mu})S_t dt + \sqrt{V_t}S_t \left[ \rho dW_t^{(1)} + \sqrt{1 - \rho^2} dW_t^{(2)} \right] + (\xi^s - 1)dN_t \quad (30)$$

$$dV_t = \kappa(\theta - V_t)dt + \sigma_v \sqrt{V_t} dW_t^{(1)} \quad (31)$$

where  $N_t$  is a Poisson process with constant intensity  $\lambda$ , and  $\xi^s$  is the relative jump size in the stock price. In particular, when a jump occurs at time  $t$ , then  $S_{t+} = S_{t-}\xi^s$ . The distribution of  $\xi^s$  is lognormal with mean  $\mu_s$  and variance  $\sigma_s^2$ . The parameters  $\mu_s$  and  $\bar{\mu}$  are related to each other by the equation:

$$\mu_s = \log(1 + \bar{\mu}) - \frac{1}{2}\sigma_s^2$$

Table 5: Simulation Results for a European Call Option using Conditional Monte Carlo

a. Simulation with the Exact Method

| No of Simulation Trials | RMS Error | Computing Time (sec.) |
|-------------------------|-----------|-----------------------|
| 10000                   | 0.0803    | 3.8                   |
| 40000                   | 0.0398    | 15.1                  |
| 160000                  | 0.0199    | 60.1                  |
| 640000                  | 0.0100    | 239.7                 |
| 2560000                 | 0.0050    | 958.9                 |
| 10240000                | 0.0025    | 3835.7                |

b. Simulation with the Euler discretization

| No of Simulation Trials | No of Time Steps | Bias   | Standard Error | RMS Error | Computing Time (sec.) |
|-------------------------|------------------|--------|----------------|-----------|-----------------------|
| 10000                   | 100              | 2.1046 | 0.1002         | 2.1070    | 0.1                   |
| 40000                   | 200              | 1.5870 | 0.0469         | 1.5877    | 1.0                   |
| 160000                  | 400              | 1.2025 | 0.0227         | 1.2027    | 8.4                   |
| 640000                  | 800              | 0.9154 | 0.0110         | 0.9155    | 64.2                  |
| 2560000                 | 1600             | 0.7022 | 0.0054         | 0.7022    | 512.1                 |
| 10240000                | 3200             | 0.5391 | 0.0026         | 0.5391    | 4086.5                |

Option parameters:  $S = 100, K = 100, V_0 = 0.09, \kappa = 2.00, \theta = 0.09, \sigma_v = 1.00, \rho = -0.30, r = 5.00\%, T = 5.0$  years, true option price=34.9998.

and only one of them needs to be specified.

To sample from the final stock price, the jump part and the diffusion part can be simulated separately. In other words, we can first simulate the diffusion part of the stock price, and then multiply by the realized jump sizes. This makes it possible to extend the exact simulation method to simulate under the SVJ model.

To generate a sample for the derivative payoff which has maturity  $T$ , we use the following procedure:

Exact Simulation Algorithm for the SVJ Model:

1. Disregard the jump part, and simulate the stock price  $S_T$  as in the SV model using the exact simulation method described in Section 3.
2. Generate a Poisson random variable with mean  $(\lambda T)$  to determine the number of jumps occurred in the time horizon, call this number  $J$ .
3. Generate independent jump sizes  $\xi_i^s$ , for  $i = 1, \dots, J$ , from the lognormal distribution with mean  $\mu_s$  and variance  $\sigma_s^2$ .
4. Find the adjusted final stock price  $\tilde{S}_T$  by multiplying the price from step 1 with the jump sizes, i.e., set  $\tilde{S}_T = S_T \prod_{i=1}^{i=J} \xi_i^s$ .
5. Compute the derivative payoff using  $\tilde{S}_T$ .

Table 6 gives numerical results for a European call option using exact simulation and Euler discretization methods. The model parameters are taken from Duffie et al. [13], these are fitted parameters for S&P 500 on a particular day. The bias of the Euler discretization method is relatively small for the set of parameters used. Therefore, we choose the number of time steps,  $M$ , to be equal to  $0.1\sqrt{N}$ , where  $N$  is the number of simulation trials. The bias column is estimated using 500 million simulation runs. Figure 5 shows the convergence of the RMS errors for the two simulation methods used. The exact method has a better RMS error than the Euler discretization method except only when less than one second of computation time is used.

## 6.2 Simulation Under the SVCJ Model

The SVCJ model introduced in Duffie et al. [13] is similar to the SVJ model, but it also includes jumps in the variance process. The governing equations are:

$$dS_t = (r - \lambda\bar{\mu})S_t dt + \sqrt{V_t}S_t \left[ \rho dW_t^{(1)} + \sqrt{1 - \rho^2} dW_t^{(2)} \right] + (\xi^s - 1)dN_t \quad (32)$$

$$dV_t = \kappa(\theta - V_t)dt + \sigma_v \sqrt{V_t} dW_t^{(1)} + \xi^v dN_t \quad (33)$$

where  $N_t$  is again a Poisson process with constant intensity  $\lambda$ ,  $\xi^s$  is the relative jump size of the stock price, and  $\xi^v$  is the jump size of the variance. The jumps in stock price and the variance occur concurrently, and their magnitudes have a correlation determined by the parameter  $\rho$ . The distribution

Table 6: Simulation Results under the SVJ model for a European Call Option

a. Simulation with the Exact Method

| No of Simulation Trials | RMS Error | Computing Time (sec.) |
|-------------------------|-----------|-----------------------|
| 10000                   | 0.2232    | 0.6                   |
| 40000                   | 0.1124    | 2.3                   |
| 160000                  | 0.0560    | 9.1                   |
| 640000                  | 0.0280    | 36.2                  |
| 2560000                 | 0.0140    | 144.7                 |
| 10240000                | 0.0070    | 579.4                 |

b. Simulation with the Euler discretization

| No of Simulation Trials | No of Time Steps | Bias   | Standard Error | RMS Error | Computing Time (sec.) |
|-------------------------|------------------|--------|----------------|-----------|-----------------------|
| 10000                   | 10               | 0.5902 | 0.2480         | 0.6402    | 0.1                   |
| 40000                   | 20               | 0.2834 | 0.1169         | 0.3065    | 0.3                   |
| 160000                  | 40               | 0.1322 | 0.0575         | 0.1442    | 1.9                   |
| 640000                  | 80               | 0.0557 | 0.0283         | 0.0625    | 12.6                  |
| 2560000                 | 160              | 0.0196 | 0.0141         | 0.0241    | 97.7                  |
| 10240000                | 320              | 0.0043 | 0.0070         | 0.0082    | 776.1                 |

Option parameters:  $S = 100$ ,  $K = 100$ ,  $V_0 = 0.008836$ ,  $\kappa = 3.99$ ,  $\theta = 0.014$ ,  $\sigma_v = 0.27$ ,  $\rho = -0.79$ ,  $\lambda = 0.11$ ,  $\bar{\mu} = -0.12$ ,  $\sigma_s = 0.15$ ,  $r = 3.19\%$ ,  $T = 5.0$  years, true option price=20.1642.

of  $\xi^v$  is exponential with mean  $\mu_v$ . Given  $\xi^v$ ,  $\xi^s$  is lognormally distributed with mean  $(\mu_s + \rho_J \xi^v)$  and variance  $\sigma_s^2$ . The parameters  $\mu_s$  and  $\bar{\mu}$  are related to each other by the equation:

$$\mu_s = \log [(1 + \bar{\mu})(1 - \rho_J \mu_v)] - \frac{1}{2} \sigma_s^2$$

and only one of them needs to be specified.

The jumps in variance prevent us from using the same method as in SVJ model but it is still possible to extend the method to generate exact simulation estimators under the SVCJ model. We need to divide the simulation according to the jump occurrences and simulate the values of variance and stock price at each jump time.

To generate a sample for the derivative payoff which has maturity  $T$ , we use the procedure described below. We denote the initial stock price by  $S_0$ , the initial variance value by  $V_0$ , and the current time by  $t_0$ :

Exact Simulation Algorithm for the SVCJ Model:

1. Simulate a Poisson process with arrival rate  $\lambda$  and determine the time of the next jump, denote this time by  $t_j$ . If  $t_j > T$  then set  $t_j = T$ .
2. Disregard the jump part, and simulate the variance value  $V_{t_j}$  and stock price  $S_{t_j}$  using the exact simulation method described in Section 3 using a time interval of  $\Delta t = t_j - t_0$ .
3. If  $t_j = T$  then set  $S_T = S_{t_j}$  and go to step 6. Otherwise generate  $\xi^v$  by sampling from an exponential distribution with mean  $\mu_v$ . Update the variance value by setting  $\tilde{V}_{t_j} = V_{t_j} + \xi^v$ .
4. Generate  $\xi^s$  by sampling from a lognormal distribution with mean  $(\mu_s + \rho_J \xi^v)$  and variance  $\sigma_s^2$ . Update the stock price by setting  $\tilde{S}_{t_j} = S_{t_j} \xi^s$ .
5. Set  $S_0 = \tilde{S}_{t_j}$ ,  $V_0 = \tilde{V}_{t_j}$ ,  $t_0 = t_j$  and go to step 1.
6. Compute the derivative payoff using  $S_T$ .

Dividing the simulation horizon and using more time steps affects the computation speed, therefore the simulation of SVCJ model is slower than SV or SVJ models. If, for example, the arrival rate of jumps is one per year, then the simulation of a 5 year option will be about 5 times slower. Some of the advantage of simulating the stock price over a long horizon is lost, but still, the method is practical and provides unbiased simulation estimators of derivative prices.

The Fourier inversion step in the exact simulation method may run into some problems when the simulation interval is too small. Such cases may occur in the simulation of SVCJ model when two jumps are very close to each other or when a jump is too close to maturity. If the simulation interval is less than 0.01 year, instead of using the exact simulation method, we use Euler discretization with 100 time steps. This introduces negligible bias into simulation estimate and solves the complexities for small time intervals.

Table 7 gives some numerical results for simulation under the SVCJ model with the exact and Euler discretization methods. The model parameters are again the fitted parameters for S&P 500 and taken from Duffie at al. [13]. For the Euler discretization, we choose the number of time steps,  $M$ , to be equal to  $0.1\sqrt{N}$ , where  $N$  is the number of simulation trials. The bias column is estimated using 500 million simulation runs. Figure 6 shows the convergence of the RMS errors for both methods. In this case, the point where the exact method starts dominating Euler discretization is closer to the right of the graph. This is both because the bias in the Euler discretization is relatively small for this set of parameters,

and also because the jumps increase the computation time of the exact simulation method.

Table 7: Simulation Results under the SVCJ model for a European Call Option

a. Simulation with the Exact Method

| No of Simulation Trials | RMS Error | Computing Time (sec.) |
|-------------------------|-----------|-----------------------|
| 10000                   | 0.0720    | 1.5                   |
| 40000                   | 0.0369    | 5.8                   |
| 160000                  | 0.0184    | 23.3                  |
| 640000                  | 0.0092    | 93.5                  |
| 2560000                 | 0.0046    | 373.0                 |
| 10240000                | 0.0023    | 1491.3                |
| 40960000                | 0.0011    | 5967.5                |

b. Simulation with the Euler discretization

| No of Simulation Trials | No of Time Steps | Bias   | Standard Error | RMS Error | Computing Time (sec.) |
|-------------------------|------------------|--------|----------------|-----------|-----------------------|
| 10000                   | 10               | 0.0148 | 0.0729         | 0.0744    | 0.1                   |
| 40000                   | 20               | 0.0090 | 0.0367         | 0.0378    | 0.2                   |
| 160000                  | 40               | 0.0050 | 0.0183         | 0.0190    | 1.6                   |
| 640000                  | 80               | 0.0024 | 0.0092         | 0.0095    | 13.3                  |
| 2560000                 | 160              | 0.0014 | 0.0046         | 0.0048    | 98.8                  |
| 10240000                | 320              | 0.0012 | 0.0023         | 0.0026    | 784.4                 |
| 40960000                | 640              | 0.0008 | 0.0011         | 0.0014    | 6254.7                |

Option parameters:  $S = 100$ ,  $K = 100$ ,  $V_0 = 0.007569$ ,  $\kappa = 3.46$ ,  $\theta = 0.008$ ,  $\sigma_v = 0.14$ ,  $\rho = -0.82$ ,  $\lambda = 0.47$ ,  $\bar{\mu} = -0.1$ ,  $\sigma_s = 0.0001$ ,  $\mu_v = 0.05$ ,  $\rho_J = -0.38$ ,  $r = 3.19\%$ ,  $T = 1.0$  year, true option price=6.8619.

## 7 An Application: Pricing of Forward Start Options

The exact simulation method of this paper can be applied to the pricing of many exotic options. To illustrate this point, in this section we consider the pricing of forward start options. These are options whose strike is set at a future date. In particular, if  $T_1$  is the time when strike is set,  $T_2$  is the option expiration,  $S_i$  is the stock price at time  $T_i$ , and  $k$  is the constant that determines the strike, then the forward start option payoff at time  $T_2$  is given by  $(S_2 - kS_1)^+$ . For example, if  $k = 1$ , then at time  $T_1$ , the option becomes an at-the-money option with expiration  $T_2$ . Kruse and Nögel [23] recently developed an expression for this option under the SV model by integrating the pricing formula with the conditional density of the variance value at time  $T_1$ . However, numerical evaluation of their expression is quite complicated since it includes another level of integration to already complex integrals in the Heston formula. Therefore, simulation may be considered as a more practical alternative for finding




Figure 5: Convergence of the RMS Errors of the Two Methods for the Option in Table 6 (SVJ Model)




Figure 6: Convergence of the RMS Errors of the Two Methods for the Option in Table 7 (SVCJ Model)

prices and sensitivities of this option. Also, for the SVJ and SVCJ models, no similar expressions are available and so other numerical methods are necessary for pricing forward start options under these models.

It is straightforward to price a forward start option using simulation: for each path, simulate the stock price values at  $T_1$  and  $T_2$ , and then evaluate the payoff function using these two values. However, we can use an alternative method for simulating a forward start option that will allow us to get more efficient estimators. Note that at time  $T_1$ , we know the stock price, strike and the expiration of the option. Therefore the option price at time  $T_1$  can be written using closed form formulas. Let  $C_E(S, K, T, V)$  denote the price of a European call option with initial stock price  $S$ , strike  $K$ , time to expiration  $T$ , and initial variance  $V$ . With this notation, at time  $T_1$ , the holder of the forward start option has a call option with value  $C_E(S_1, kS_1, T_2 - T_1, V_1)$ , where  $V_1$  is the variance value at  $T_1$ . Note also that, the option price is linearly homogenous with respect to the stock price and the strike, i.e., we can write  $C_E(S, kS, T, V) = S C_E(1, k, T, V)$ . Using this expression, and following the above arguments, the price of a forward start option can be written as:

$$C_{FW} = E[e^{-rT_1} S_1 C_E(1, k, T_2 - T_1, V_1)] \quad (34)$$

Thus, we can price forward start options by simulating the stock price and the variance values at time  $T_1$ , and then using the European option pricing formula to evaluate the option payoff. Using the analytical formula in expression (34) eliminates the variance that occurs due to the uncertainty after  $T_1$ . Therefore, this estimator is expected to give a better estimate than plain Monte Carlo simulation.

In Table 8, we give simulation results for a forward start option under each model. The model parameters are taken from Duffie et al. [13], and are the same as given in the previous sections. The price estimator given in (34) is denoted as ‘Formula Simulation Price’, and the plain Monte Carlo estimate that simulates stock prices at  $T_1$  and  $T_2$  is denoted as ‘Plain Simulation Price’. The computation time per simulation path is different for the two methods. We adjust the number of simulation trials for the ‘Formula Simulation’ method such that it takes roughly the same amount of time as the ‘Plain Simulation’ method. Both simulation estimators are unbiased since we use the exact simulation algorithm, but combining the algorithm with the analytical European call formula results in significant variance reduction.

Table 8: Simulation Estimates for a Forward Start Option

|                          | MODEL  |        |        |
|--------------------------|--------|--------|--------|
|                          | SV     | SVJ    | SVCJ   |
| Plain Simulation Price   | 7.0213 | 7.0352 | 7.1376 |
| standard error           | 0.0778 | 0.0777 | 0.0798 |
| No of Simulation Trials  | 10,000 | 10,000 | 10,000 |
| Computing Time (sec.)    | 6.1    | 1.7    | 2.9    |
| Formula Simulation Price | 6.9708 | 6.8978 | 7.0593 |
| standard error           | 0.0088 | 0.0149 | 0.0136 |
| No of Simulation Trials  | 8,600  | 2,800  | 3,300  |
| Computing Time (sec.)    | 6.1    | 1.7    | 2.9    |

Option Parameters:  $S_0=100$ ,  $k=1$ ,  $T_1=1$  year,  $T_2=2$  years. SV parameters are the same as in Table 1; SVJ parameters are the same as in Table 6; SVCJ parameters are the same as in Table 7.

## 8 Conclusion

In this paper we propose a simulation method for exact simulation of stock price and variance under Heston's stochastic volatility and some other affine jump diffusion models. Our method is based on Fourier inversion techniques and conditioning arguments. The proposed simulation method recovers the error convergence rate of  $O(s^{-\frac{1}{2}})$  for unbiased Monte Carlo simulation, where  $s$  is the total computation time. Using the more conventional Euler discretization method leads to  $O(s^{-\frac{1}{3}})$  or slower convergence, depending on the convergence rate of the bias.

The proposed simulation method may be used to generate unbiased price estimators for path-independent derivatives, and also for some path-dependent derivatives with mild path-dependence. By mild path-dependence, we mean that the derivative payoff should depend on the stock price and variance values at few points during the life of the derivative. Bermudan options and American call options with discrete dividends are some examples. In the case of path-independent derivatives, any type of payoff that depends on the final stock price and variance can be handled. Also, under the SV model, it is possible to generate unbiased estimators of the Greeks of path-independent derivatives using the conditional Monte Carlo method.

The exact simulation method has several advantages over the discretization methods. The bias for the discretization methods may be large and many time steps may be needed to reduce the bias to an

acceptable level, which increases the computation time. Also, since the bias is unknown, the standard error may be a poor estimate of the actual error, and valid confidence intervals are not available. Finally, it is difficult to determine the optimal tradeoff between the time steps and simulation trials since the convergence rate of the bias is unknown.

We verified the faster convergence rate of our exact simulation methods through numerical examples. We demonstrated that they achieve a smaller RMS error than the Euler discretization under SV and SVJ models, except when a very small amount of computation time is used. The exact simulation method is relatively slower for simulation under the SVCJ model, but it is still more efficient than the Euler discretization when high accuracy is needed. We have also shown that the conditional Monte Carlo methods of Willard [31] can be used to improve the efficiency of the exact method for simulation of certain path-independent options under the SV model.

## 9 Appendix: Derivation of the Laplace Transform

In this appendix we derive the Laplace Transform for the integral of  $\int_u^t V(s)ds$  given  $V(u)$  and  $V(t)$ . We follow the approach of Chesney et al. [10] to get a square-root process with volatility parameter  $\sigma = 2$  and then use the results of Pitman and Yor [27] to write the Laplace Transform.

$V(t)$  follows the following square-root process:

$$V(t) = V(0) + \int_0^t \kappa[\theta - V(s)]ds + \int_0^t \sigma \sqrt{V(s)}dW_s. \quad (35)$$

Now,

$$\begin{aligned} \int_0^t \sigma \sqrt{V(s)}dW_s &= 2 \int_0^t \sqrt{V(s)} \frac{\sigma}{2} dW_s \\ &\cong 2 \int_0^t \sqrt{V(s)} dW_{\sigma^2 s/4} \quad (\text{in law}) \\ &= 2 \int_0^t \sqrt{V\left(\frac{4}{\sigma^2} \frac{\sigma^2}{4} s\right)} dW_{\sigma^2 s/4}. \end{aligned} \quad (36)$$

The second equality in (36) follows from the scaling property of Brownian motion. Define  $u = \sigma^2 s/4$ ,

so we have  $du = (\sigma^2/4)ds$ . We can write (35) as

$$V\left(\frac{4}{\sigma^2}\frac{\sigma^2}{4}t\right) = V(0) + \frac{4}{\sigma^2} \int_0^{\sigma^2 t/4} \kappa \left[ \theta - V\left(\frac{4}{\sigma^2}u\right) \right] du + 2 \int_0^{\sigma^2 t/4} \sqrt{V\left(\frac{4}{\sigma^2}u\right)} dW_u.$$

Defining the process  $\rho(u) = V(4u/\sigma^2)$ , we obtain

$$\rho(\sigma^2 t/4) = \rho(0) + \frac{4}{\sigma^2} \int_0^{\sigma^2 t/4} \kappa[\theta - \rho(u)] du + 2 \int_0^{\sigma^2 t/4} \sqrt{\rho(u)} dW_u. \quad (37)$$

Setting  $n = 4\kappa\theta/\sigma^2$  and  $j = -2\kappa/\sigma^2$ , we get

$$\rho(t) = \rho(0) + \int_0^t [2j\rho(u) + n] du + 2 \int_0^t \sqrt{\rho(u)} dW_u. \quad (38)$$

The infinitesimal generator of this process is  $2xD^2 + (2jx + n)D$  where  $D = \frac{d}{dx}$ . Pitman and Yor [27] give the following formula for the squared Bessel process  $X(s)$  which has generator  $2xD^2 + nD$ :

$$\begin{aligned} \tilde{E} \left[ \exp \left\{ -\frac{b^2}{2} \int_0^t X(s) ds \right\} \middle| X(0) = x, X(t) = y \right] &= \left( \frac{bt}{\sinh bt} \right) \exp \left\{ \frac{x+y}{2t} (1 - bt \coth bt) \right\} \\ &\times I_\nu \left( \frac{\sqrt{xy}b}{\sinh bt} \right) / I_\nu \left( \frac{\sqrt{xy}}{t} \right). \end{aligned} \quad (39)$$

where  $\nu = n/2 - 1$ ,  $I_\nu(\cdot)$  is the modified Bessel function of the first kind, and  $\tilde{E}$  denotes the expectation taken with respect to the law of the squared Bessel process. To be able to use this formula for the process in (38), we need to apply a change of law formula to eliminate the random component of the drift term and get a process with  $j = 0$ .

We now write the Laplace transform we want to derive:

$$\begin{aligned}
E \left[ \exp \left\{ -a \int_u^t V(s) ds \right\} \middle| V(u), V(t) \right] &= E \left[ \exp \left\{ -a \int_u^t \rho\left(\frac{\sigma^2 s}{4}\right) ds \right\} \middle| \rho\left(\frac{\sigma^2 u}{4}\right), \rho\left(\frac{\sigma^2 t}{4}\right) \right] \\
&= E \left[ \exp \left\{ -\frac{4a}{\sigma^2} \int_{\sigma^2 u/4}^{\sigma^2 t/4} \rho(s) ds \right\} \middle| \rho\left(\frac{\sigma^2 u}{4}\right), \rho\left(\frac{\sigma^2 t}{4}\right) \right] \\
&= \tilde{E} \left[ \exp \left\{ -\left(\frac{4a}{\sigma^2} + \frac{j^2}{2}\right) \int_{\sigma^2 u/4}^{\sigma^2 t/4} \rho(s) ds \right\} \middle| \rho\left(\frac{\sigma^2 u}{4}\right), \rho\left(\frac{\sigma^2 t}{4}\right) \right] / \\
&\tilde{E} \left[ \exp \left\{ -\frac{j^2}{2} \int_{\sigma^2 u/4}^{\sigma^2 t/4} \rho(s) ds \right\} \middle| \rho\left(\frac{\sigma^2 u}{4}\right), \rho\left(\frac{\sigma^2 t}{4}\right) \right].
\end{aligned}$$

The last equality follows from the change of law formula(6.d) of Pitman and Yor [27]. The expectations in the numerator and the denominator are calculated using the formula (39).

After arranging the terms and changing back to  $V(\cdot)$  domain, we get the Laplace transform as:

$$\begin{aligned}
E \left[ \exp \left\{ -a \int_u^t V(s) ds \right\} \middle| V(u), V(t) \right] &= \frac{\gamma(a) e^{\frac{-1}{2}(\gamma(a)-\kappa)(t-u)} (1 - e^{-\kappa(t-u)})}{\kappa(1 - e^{-\gamma(a)(t-u)})} \\
&\times \exp \left\{ \frac{V(u) + V(t)}{\sigma^2} \left[ \frac{\kappa(1 + e^{-\kappa(t-u)})}{1 - e^{-\kappa(t-u)}} - \frac{\gamma(a)(1 + e^{-\gamma(a)(t-u)})}{1 - e^{-\gamma(a)(t-u)}} \right] \right\} \\
&\times \frac{I_{0.5n-1} \left[ \frac{4\gamma(a)\sqrt{V(u)V(t)}}{\sigma^2} \frac{e^{-0.5\gamma(a)(t-u)}}{(1 - e^{-\gamma(a)(t-u)})} \right]}{I_{0.5n-1} \left[ \frac{4\kappa\sqrt{V(u)V(t)}}{\sigma^2} \frac{e^{-0.5\kappa(t-u)}}{(1 - e^{-\kappa(t-u)})} \right]} \tag{40}
\end{aligned}$$

where  $\gamma(a) = \sqrt{\kappa^2 + 2\sigma^2 a}$ .

## References

- [1] Abate, J., and Whitt, W., (1992), The Fourier-Series Method for Inverting Transforms of Probability Distributions, *Queueing Systems*, Vol.10, No.1, 5-88.
- [2] Abramowitz, M., and Stegun, I.A., (1972), *Handbook of Mathematical Functions with Formulas*,

*Graphs, and Mathematical Tables*, National Bureau of Standards, Washington D.C.

- [3] Amos, D. E., (1986), Algorithm 644, A Portable Package for Bessel Functions of a Complex Argument and Nonnegative Order, *ACM Trans. Math. Softw.*, Vol.12, No.3, 265-273.
- [4] Bakshi, G., Cao, C. and Chen, Z., (1997), Empirical Performance of Alternative Option Pricing Models, *Journal of Finance*, Vol.52, 2003-2049.
- [5] Bally, V., and Talay, D., (1996), The Law of the Euler Scheme for Stochastic Differential Equations I: Convergence Rate of the Distribution Function, *Probability Theory and Related Fields*, Vol.104, 43-60.
- [6] Bates, D., (1996), Jumps and Stochastic Volatility: Exchange Rate Processes Implicit in Deutsche Mark Options, *Review of Financial Studies*, Vol.9, 69-107.
- [7] Broadie, M., and Glasserman, P., (1996), Estimating Security Price Derivatives Using Simulation, *Management Science*, Vol.42, No.2, 269-285.
- [8] Broadie, M., and Kaya, Ö., (2004), Exact Simulation of Option Greeks under Stochastic Volatility and Jump Diffusion Models, in *Proceedings of the 2004 Winter Simulation Conference*, eds: R.G. Ingalls, M.D. Rossetti, J.S. Smith, and B.A. Peters, The Society for Computer Simulation.
- [9] Chernov, M., Ghysels, E., Gallant, A. and Tauchen, G., (2003), Alternative Models for Stock Price Dynamics, *Journal of Econometrics*, Vol.116, 225-257.
- [10] Chesney, M., Elliot, R.J. and Gibson, R., (1993), Analytical Solutions for the Pricing of American Bond and Yield Options, *Mathematical Finance*, Vol.3, No.3, 277-294.
- [11] Cox, J.C., Ingersoll, J.E., and Ross, S.A., (1985), A theory of the term structure of interest rates, *Econometrica*, Vol.53, No.2, 385-407.
- [12] Duffie, D. and Glynn, P., (1995), Efficient Monte Carlo Estimation of Security Prices, *The Annals of Applied Probability*, Vol.4, No.5, 897-905.
- [13] Duffie, D., Singleton, K. and Pan, J., (2000), Transform Analysis and Asset Pricing for Affine Jump-Diffusions, *Econometrica*, Vol.68, 1343-1376.

- [14] Feller, W., (1971), *An Introduction to Probability Theory and Its Applications, Volume II*, second edition, John Wiley & Sons, New York.
- [15] Eraker, B., Johannes, M. and Polson, N., (2003), The Impact of Jumps in Volatility and Returns, *Journal of Finance*, Vol.58, No.3, 1269-1300.
- [16] Fishman, G.S., (1996), *Monte Carlo: Concepts, Algorithms, and Applications*, Springer-Verlag, New York.
- [17] Glasserman, P., (2003), *Monte Carlo Methods in Financial Engineering*, Springer-Verlag, New York.
- [18] Heath, D. and Platen, E., (2002), A Variance Reduction Technique based on Integral Representations, *Quantitative Finance*, Vol.2, No.5, 362-369.
- [19] Heston, S., (1993), A Closed-Form Solution of Options with Stochastic Volatility with Applications to Bond and Currency Options, *The Review of Financial Studies*, Vol.6, No.2, 327-343.
- [20] Hull, J., and White, A., (1987), The Pricing of Options on Assets with Stochastic Volatilities, *Journal of Finance*, Vol.XLII, No.2, 281-300.
- [21] Johnson, N.L., Kotz, S., and Balakrishnan, N., (1994), *Continuous Univariate Distributions, Volume 2*, Second Edition, Wiley, New York.
- [22] Kloeden, P., and Platen, E., (1992), *Numerical Solution of Stochastic Differential Equations*, Springer-Verlag, New York.
- [23] Kruse, S., and Nögel, U., (2004), On the Pricing of Forward Starting Options in Heston's Model on Stochastic Volatility, Working Paper, Fraunhofer-ITWM, Kaiserslautern.
- [24] Matsumoto M., and Nishimura T., (1998), Mersenne Twister: A 623-dimensionally equidistributed uniform pseudorandom number generator, *ACM Transactions on Modeling and Computer Simulation*, Vol.8, No.1, 3-30.
- [25] Moro, B., (1995), The Full Monte, *RISK*, Vol.8, No.2, 57-58.

- [26] Pan, J., (2002), The Jump-Risk Premia Implicit in Options: Evidence from an Integrated Time-Series Study, *Journal of Financial Economics*, Vol.63, 3-50.
- [27] Pitman, J., and Yor, M., (1982), A Decomposition of Bessel Bridges, *Z. Wahrscheinlichkeitstheorie verw. Gebiete*, Vol. 59, 425-457.
- [28] Romano, M., and Touzi, N., (1997), Contingent Claims and Market Completeness in a Stochastic Volatility Model, *Mathematical Finance*, Vol.7, No.4, 399-410.
- [29] Scott, L.O., (1996), Simulating a Multi-Factor Term Structure Model Over Relatively Long Discrete Time Periods, in *Proceedings of the IAFE First Annual Computational Finance Conference*, Graduate School of Business, Stanford University.
- [30] Talay, D., and Tubaro, L., (1990), Expansion of the Global Error for Numerical Schemes Solving Stochastic Differential Equations, *Stochastic Analysis and Applications*, Vol.8, 94-120.
- [31] Willard, G.A., (1997), Calculating Prices and Sensitivities for Path-Independent Derivative Securities in MultiFactor Models, *The Journal of Derivatives*, Vol.5, No.1, 45-61.